



TITLE:

Topics on Free Boundary Problems for Ideal Fluids (Mathematical Analysis in Fluid and Gas Dynamics)

AUTHOR(S):

Tani, Atushi

CITATION:

Tani, Atushi. Topics on Free Boundary Problems for Ideal Fluids (Mathematical Analysis in Fluid and Gas Dynamics). 数理解析研究所講究録 2004, 1353: 35-48

ISSUE DATE:

2004-01

URL:

<http://hdl.handle.net/2433/25145>

RIGHT:

Topics on Free Boundary Problems for Ideal Fluids

慶應義塾大学・理工学部 谷 温之 (Atusi TANI)

Department of Mathematics, Keio University

This article reviews the free boundary problems for the motion of an incompressible ideal fluid. These problems are typically classified into the following three types according to the geometrical configurations of fluid domain:

- [I] Vortical water waves, i.e., fluid motion in a domain of infinite extent bounded by the upper free surface and the lower bottom of finite or infinite depth (in this case a dominant external force is due to a gravitation downward vertically),
- [II] Circulating fluid around a celestial body, i.e., fluid motion around a rigid body with a compact free surface (in this case a dominant external force is due to a gravitation of the celestial body),
- [III] Gaseous stars, i.e., fluid motion in a domain bounded by the free surface (in this case a dominant external force is due to a self-gravitational force).

In general, as the vector fields of external forces we should take not only the potential but the general form. Besides the forces mentioned above the effect of the surface tension is taken into account.

We have a long history and a lot of works discussing on these problems, however concerning the most fundamental study of the wellposedness of these problems there are not so many works. In this article we are interested in just this wellposedness.

We begin with a classical description of the problem.

At time $t > 0$ let $\Omega(t)$ be a domain in \mathbb{R}^N ($N = 2, 3$) occupied by the fluid, which is bounded by a bottom Γ_b and a free surface $\Gamma_s(t)$:

$$\begin{aligned}\Gamma_b &= \{\mathbf{x} \in \mathbb{R}^N \mid F_b(\mathbf{x}) = 0\}, \\ \Gamma_s(t) &= \{\mathbf{x} \in \mathbb{R}^N \mid F_s(\mathbf{x}, t) = 0\}.\end{aligned}$$

We always assume that $\Gamma_b \cap \Gamma_s(t) = \emptyset$ for any $t \geq 0$. Corresponding to problems [I], [II] and [III], we usually consider the fluid motion in the domains given by

$$F_b(\mathbf{x}) \equiv x_N - b(\mathbf{x}'), \quad F_s(\mathbf{x}, t) \equiv x_N - \eta(\mathbf{x}', t) \quad (\mathbf{x} = (\mathbf{x}', x_N)),$$

$$F_b(\mathbf{x}) \equiv |\mathbf{x}| - b(\omega), \quad F_s(\mathbf{x}, t) \equiv |\mathbf{x}| - \eta(\omega, t) \quad (\omega \in S^{N-1}),$$

$$\Gamma_b = \emptyset, \quad F_s(\mathbf{x}, t) \equiv |\mathbf{x}| - \eta(\omega, t) \quad (\omega \in S^{N-1}),$$

respectively, where S^{N-1} is an $(N-1)$ -dimensional sphere.

The motion of an incompressible ideal fluid is described in the Eulerian coordinates by

$$(1) \quad \rho \frac{D}{Dt} \mathbf{v} + \nabla_x p = \rho \mathbf{f}, \quad \operatorname{div}_x \mathbf{v} = 0 \quad \text{in } \Omega(t), \quad t > 0,$$

where $\mathbf{v} = \mathbf{v}(\mathbf{x}, t)$ is the velocity vector field, $p = p(\mathbf{x}, t)$ is the pressure, $\mathbf{f} = \mathbf{f}(\mathbf{x}, t)$ is a vector field of exterior mass forces, ρ is the constant density of the fluid and $D/Dt = \partial/\partial t + \mathbf{v} \cdot \nabla_x$ is a material derivative.

Note that correspondent to problems [I], [II] and [III], the suitable forms of the external forces are considered as

$$\begin{aligned} \mathbf{f}(\mathbf{x}, t) &= -\rho g \nabla \Phi(\mathbf{x}, t), \quad \Phi(\mathbf{x}, t) = x_{\perp}, \\ \mathbf{f}(\mathbf{x}, t) &= \rho M g \nabla \Phi(\mathbf{x}, t), \quad \Phi(\mathbf{x}, t) = \frac{1}{|\mathbf{x}|}, \\ \mathbf{f}(\mathbf{x}, t) &= 4\pi \rho g \nabla \Phi(\mathbf{x}, t), \quad \Phi(\mathbf{x}, t) = \int_{\Omega(t)} \frac{\rho}{|\mathbf{x} - \mathbf{y}|} dy, \end{aligned}$$

where g is a gravitational constant, x_{\perp} is a perpendicular component of \mathbf{x} and M is a mass of a celestial body. In problem [II] we take the barycenter of the celestial body as an origin of the coordinate system.

The boundary conditions on $\Gamma_s(t)$ are

$$(2) \quad \begin{cases} \frac{D}{Dt} F = 0 & (\text{kinematic condition}), \\ p = p_e + 2\sigma H & (\text{dynamic condition}), \end{cases}$$

and on Γ_b

$$(3) \quad (\mathbf{v} - \mathbf{v}_b) \cdot \mathbf{n}_b = 0.$$

Here p_e is an atmosphere pressure, $\sigma (\geq 0)$ a coefficient of surface tension, H a mean curvature ($H > 0$ if Γ_t is convex outside the fluid region), \mathbf{v}_b velocity of Γ_b and \mathbf{n}_b a normal vector to Γ_b .

Initial conditions are

$$(4) \quad \begin{cases} \Omega(0) = \Omega \quad (\Gamma_s(0) = \Gamma_s), \\ \mathbf{v}|_{t=0} = \mathbf{v}_0(\mathbf{x}), \quad \operatorname{div} \mathbf{v}_0 = 0, \quad \mathbf{x} \in \Omega. \end{cases}$$

Our aim is to find a solution $(\Omega(t), \mathbf{v}(\mathbf{x}, t), p(\mathbf{x}, t))$ for $t > 0$ to problem (1) – (4).

It is convenient to write the problem in the Lagrangean coordinates ξ :

$$(5) \quad \frac{d}{dt} \mathbf{x} = \mathbf{v}(\mathbf{x}, t), \quad \mathbf{x}|_{t=0} = \xi,$$

which can be solved by the formula

$$(5)' \quad \mathbf{x} = \xi + \int_0^t \mathbf{u}(\xi, \tau) d\tau, \quad \mathbf{u}(\xi, t) = \mathbf{v}(\mathbf{x}(\xi, t), t).$$

Let $\mathcal{G}(\xi, t) = G(\mathbf{x}(\xi, t), t)$. Then from (5) it follows that

$$\frac{\partial}{\partial t} \mathcal{G} = \frac{D}{Dt} G.$$

In particular, $\mathcal{F}(\xi, t) \equiv F(\mathbf{x}, t)$ satisfies

$$\frac{\partial}{\partial t} \mathcal{F} = 0, \quad \text{hence } \Gamma_s = \{\xi \in \mathbb{R}^N \mid \mathcal{F}(\xi) = 0\}.$$

One can easily check that the mapping $\xi \mapsto \mathbf{x}$ is ont-to-one from Ω onto $\Omega(t)$, from Γ_s onto $\Gamma_s(t)$ and from Γ_b onto Γ_b . Let \mathcal{M} be a Jacobian matrix

$$\mathcal{M} = \frac{\partial(\mathbf{x})}{\partial(\xi)} = \left(\frac{\partial x_j}{\partial \xi_k} \right)_{j,k=1,2,\dots,N}.$$

Then (5) yields

$$\frac{\partial}{\partial t} \mathcal{M} = \frac{\partial(\mathbf{v})}{\partial(\mathbf{x})} \mathcal{M},$$

from which

$$(6) \quad |\mathcal{M}| \equiv \det \mathcal{M} = 1 \quad (\text{Liouville's theorem}).$$

Noting that

$$\nabla_{\mathbf{x}} = (\mathcal{M}^*)^{-1} \nabla_{\xi} \equiv \nabla_{\mathbf{u}}$$

with \mathcal{M}^* being the transposed matrix of \mathcal{M} , we transform problem (1)-(4) into the following problem in the fixed domain, which is denoted by Problem A.

Problem A. Find $(\mathbf{u}(\xi, t), p(\xi, t))$ satisfying

$$\begin{cases} \frac{\partial \mathbf{u}}{\partial t} + \frac{1}{\varrho} \nabla_{\mathbf{u}} p - \mathbf{f} = 0 & \text{in } \Omega, \quad t > 0, \\ \nabla_{\mathbf{u}} \cdot \mathbf{u} = 0 & \text{in } \Omega, \quad t > 0, \\ p = p_e + 2\sigma H & \text{on } \Gamma_s, \quad t > 0, \\ (\mathbf{v} - \mathbf{v}_b) \cdot \mathbf{n}_b = 0 & \text{on } \Gamma_b, \quad t > 0, \\ \mathbf{u}|_{t=0} = \mathbf{v}_0(\xi) & \text{on } \Omega \quad (\text{div} \mathbf{v}_0 = 0). \end{cases}$$

When $\mathbf{f} = \nabla_{\mathbf{x}} h$ and $p_e = \text{constant}$, one can deduce from Problem A to the following problem.

Problem A'. Find $(\mathbf{u}(\xi, t), q(\xi, t))$ satisfying

$$\begin{cases} \frac{\partial \mathbf{u}}{\partial t} + \frac{1}{\varrho} \nabla_{\mathbf{u}} q = 0 & \text{in } \Omega, \quad t > 0, \\ \nabla_{\mathbf{u}} \cdot \mathbf{u} = 0 & \text{in } \Omega, \quad t > 0, \\ q = -\varrho h + 2\sigma H & \text{on } \Gamma_s, \quad t > 0, \\ (\mathbf{v} - \mathbf{v}_b) \cdot \mathbf{n}_b = 0 & \text{on } \Gamma_b, \quad t > 0, \\ \mathbf{u}|_{t=0} = \mathbf{v}_0(\xi) & \text{on } \Omega \quad (\text{div} \mathbf{v}_0 = 0), \end{cases}$$

where $q = p - p_e - \rho h$.

For Problem A' in case [I] we have the following works. [?], [?], [38], [64], [65].

For Problem A' in case [I] with rotation free case we have the following works. [52], [53], [54], [55], [?], [?], [?], [94], [95]. We give another formulation.

Since operating the material derivative to equation (5) implies

$$\mathbf{x}_{tt} = \frac{D}{Dt} \mathbf{v},$$

one can derive from (1)

$$\mathcal{M}^* (\mathbf{x}_{tt} - \mathbf{f}) + \frac{1}{\rho} \nabla_{\xi} p = 0.$$

Equation (6) is indeed equivalent to equation (1)², however it is not a divergence form. Following Ovsjannikov, we derive an equivalent divergence form.

For any $\mathcal{A}_j(\mathbf{x}(\xi))$ it holds that

$$\frac{\partial}{\partial \xi_k} \mathcal{A}_j = \mathbf{x}_{\xi_k} \cdot \nabla_{\mathbf{x}} \mathcal{A}_j = \operatorname{div}_{\mathbf{x}} (\mathbf{x}_{\xi_k} \mathcal{A}_j) - \mathcal{A}_j \operatorname{div}_{\mathbf{x}} \mathbf{x}_{\xi_k},$$

and for any Jacobian matrix \mathcal{A}

$$\operatorname{div}_{\mathbf{x}} \mathbf{x}_{\xi_k} = \frac{1}{|\mathcal{M}|} \frac{\partial}{\partial \xi_k} |\mathcal{M}|.$$

If $|\mathcal{M}|$ is constant, then it holds that for any $\mathcal{A} = (\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3)$

$$\operatorname{div}_{\xi} \mathcal{A} = \operatorname{div}_{\mathbf{x}} (\mathcal{M} \mathcal{A})$$

In our case $|\mathcal{M}| = 1$. Thus $\mathcal{M} \mathcal{A} = \mathbf{x}_t = \mathbf{v}$ and (1.1)² yield

$$\operatorname{div}_{\xi} (\mathcal{M}^{-1} \mathbf{x}_t) = 0.$$

Problem B. To find $(\mathbf{x}(\xi, t), p(\xi, t))$ satisfying

$$(7) \quad \begin{cases} \mathcal{M}^* (\mathbf{x}_{tt} - \mathbf{f}) + \frac{1}{\rho} \nabla_{\xi} p = 0 & \text{in } \Omega, \ t > 0, \\ \operatorname{div}_{\xi} (\mathcal{M}^{-1} \mathbf{x}_t) = 0 \text{ or (1.6)} & \text{in } \Omega, \ t > 0, \\ p = p_e + 2\sigma H & \text{on } \Gamma_s, \ t > 0, \\ (\mathbf{x}_t - \mathbf{v}_b) \cdot \mathbf{n}_b = 0 & \text{on } \Gamma_b, \ t > 0, \\ (\mathbf{x}, \mathbf{x}_t)|_{t=0} = (\xi, \mathbf{v}_0(\xi)) & \text{on } \Omega \quad (\operatorname{div} \mathbf{v}_0 = 0). \end{cases}$$

Following Weber (1868) (see, for example [81]), we proceed further to deduce the equivalent problem to (4.1) when $\mathbf{f} = \nabla_{\mathbf{x}} h$. Since

$$\mathcal{M}^* \mathbf{x}_{tt} = \frac{\partial}{\partial t} (\mathcal{M}^* \mathbf{x}_t) - \mathcal{M}_t^* \mathbf{x}_t = \frac{\partial}{\partial t} (\mathcal{M}^* \mathbf{x}_t) - \frac{1}{2} \nabla_{\xi} |\mathbf{x}_t|^2,$$

(1.8)¹ is equivalent to

$$(8) \quad \frac{\partial}{\partial t} (\mathcal{M}^* \mathbf{x}_t) + \nabla_{\xi} \left(p - h - \frac{1}{2} |\mathbf{x}_t|^2 \right) = 0.$$

Applying the rotation operator to (8) leads to

$$\frac{\partial}{\partial t} (\text{rot}_{\xi} \mathcal{M}^* \mathbf{x}_t) = 0,$$

hence

$$\text{rot}_{\xi} \mathcal{M}^* \mathbf{x}_t = \text{rot}_{\xi} \mathbf{v}_0.$$

Then one can see that there exists φ such that

$$\mathcal{M}^* \mathbf{x}_t = \nabla_{\xi} \varphi + \mathbf{v}_0.$$

Note that this φ is generally multi-valued and single-valued if Ω is simply connected. Substituting this into (8) and integration with respect to t imply

$$\varphi_t + p = h + \frac{1}{2} |\mathbf{x}_t|^2 + \chi(t) \quad \text{for } \forall \chi(t).$$

In the following we set $\chi(t) \equiv 0$. From (7)² it follows

$$\text{div}_{\xi} (\mathcal{M}^{-1} \mathcal{M}^{*-1} (\nabla_{\xi} \varphi + \mathbf{v}_0)) = 0.$$

Finally we arrive at the equivalent problem to **Problem B**:

Problem B'. To find $(\mathbf{x}(\xi, t), \varphi(\xi, t))$ satisfying

$$(9) \quad \begin{cases} \mathcal{M}^* \mathbf{x}_t = \nabla_{\xi} \varphi + \mathbf{v}_0 & \text{in } \Omega, \quad t > 0, \\ \text{div}_{\xi} (\mathcal{M}^{-1} \mathcal{M}^{*-1} (\nabla_{\xi} \varphi + \mathbf{v}_0)) = 0 & \text{in } \Omega, \quad t > 0, \\ \varphi_t = h + \frac{1}{2} |\mathcal{M}^{*-1} (\nabla_{\xi} \varphi + \mathbf{v}_0)|^2 - p_e - 2\sigma H & \text{on } \Gamma_s, \quad t > 0, \\ (\mathcal{M}^{*-1} (\nabla_{\xi} \varphi + \mathbf{v}_0) - \mathbf{v}_b) \cdot \mathbf{n}_b = 0 & \text{on } \Gamma_b, \quad t > 0, \\ (\mathbf{x}, \varphi)|_{t=0} = (\xi, 0) & \text{on } \Omega. \end{cases}$$

We remark a little bit further. Let $\omega \equiv \text{rot}_{\mathbf{x}} \mathbf{v}$. Then (1.1)¹ yields

$$\frac{D}{Dt} \omega - \omega \cdot \nabla_{\mathbf{x}} \mathbf{v} = \text{rot}_{\mathbf{x}} \mathbf{f}.$$

When $\mathbf{f} = \nabla_{\mathbf{x}} h$, this equation becomes

$$(10) \quad \frac{D}{Dt} \omega - \omega \cdot \nabla_{\mathbf{x}} \mathbf{v} = 0.$$

Therefore

$$\omega = \mathcal{M} \text{rot}_{\xi} \mathbf{v}_0$$

follows, which is known as Lagrange-Cauchy theorem (cf. [81]). Then it is easily seen that $\text{rot}_x \mathbf{f} = 0$, $\text{rot}_x \mathbf{v}_0 = 0$ if and only if $\omega \equiv 0$.

In the case of rotation free there exist a velocity potential Φ such that $\mathbf{v} = \nabla_x \Phi$. Then $\mathcal{M}^* \mathbf{x}_t = \nabla_{\xi} \Phi$, which implies $\text{rot}_{\xi}(\mathcal{M}^* \mathbf{x}_t) = 0$. This means that $\mathcal{M}^* \mathbf{x}_t$ is a potential if and only if $\mathcal{M}^* \mathcal{M}_t$ is symmetric, i.e., $\mathcal{M}^* \mathcal{M}_t = \mathcal{M}_t^* \mathcal{M}$. For a potential flow it follows from (8) that

$$\Phi_t + p = h + \frac{1}{2} |\mathbf{x}_t|^2.$$

Then one can see that Φ is connected with φ appeared in the Weber transformation:

$$\Phi = \varphi + \Phi_0, \quad \Phi_0 = \Phi|_{t=0}.$$

In the two-dimensional case, since $\mathbf{v} = (v_1, v_2, 0)(x, x_2)$, \mathcal{M} and $\omega_3 \equiv \omega$ become

$$\mathcal{M} = \begin{pmatrix} x_{1,\xi_1} & x_{1,\xi_2} & 0 \\ x_{2,\xi_1} & x_{2,\xi_2} & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and

$$\omega = \frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2}.$$

For a potential flow (4.4) becomes

$$\omega = \omega_0(\xi_1, \xi_2) = \frac{\partial v_{0,2}}{\partial \xi_1} - \frac{\partial v_{0,1}}{\partial \xi_2}.$$

From the equation

$$\omega = \text{rot}_x = \mathcal{M} \text{rot}_{\xi} \mathcal{M}^* \mathbf{x}_t$$

it follows that

$$(11)^1 \quad x_{1,\xi_2} x_{1,\xi_1 t} - x_{1,\xi_1} x_{1,\xi_2 t} + x_{2,\xi_2} x_{2,\xi_1 t} - x_{2,\xi_1} x_{2,\xi_2 t} = \omega_0(\xi_1, \xi_2).$$

(6) implies

$$(11)^2 \quad x_{1,\xi_1} x_{2,\xi_2} - x_{1,\xi_2} x_{2,\xi_1} = 1.$$

(7)³ becomes

$$\tau \cdot \nabla_{\xi} p = \tau \cdot \nabla_{\xi} p_e + 2\sigma \tau \cdot \nabla_{\xi} H \quad \text{on } \Gamma_s,$$

where $\tau = (\tau_1, \tau_2)$ is a tangential vector to Γ_s . This, together with (7)¹, yields

$$(11)^3 \quad [x_{1,\xi_1} (x_{1,tt} - f_1) + x_{2,\xi_1} (x_{2,tt} - f_2)] \tau_1 + [x_{1,\xi_2} (x_{1,tt} - f_1) + x_{2,\xi_2} (x_{2,tt} - f_2)] \tau_2 \\ = -\frac{1}{\varrho} \tau \cdot \nabla_{\xi} p_e - \frac{2\sigma}{\varrho} \tau \cdot \nabla_{\xi} H.$$

Summing up, the two-dimensional problem for a potential flow is to find \mathbf{x} satisfying

$$\begin{cases} (11)^1, (11)^2 & \text{in } \Omega, \ t > 0, \\ (11)^3 & \text{on } \Gamma_s, \ t > 0, \\ (7)^4 & \text{on } \Gamma_b, \ t > 0, \\ (7)^5 & \text{in } \Omega \end{cases}$$

under the conditions $\operatorname{div}_{\xi} \mathbf{v}_0 = 0, \operatorname{rot}_{\xi} \mathbf{v}_0 = \omega_0$.

For Problem B' in the three dimensional and the rotation free case Bimenov proved the existence result by virtue of Nash-Moser implicit theorms [17], [18]. Using the same way, we shall be able to prove the well-posedness for the vortical case.

And Andreev [8], [9] discussed the discussed th stability for Problem B. Following his method, we shall be able to construct the solution around Gerstner's trocoidal solution.

References

- [1] Akyras, T. R., Three-dimensional long water-wave phenomena. *Ann. Rev. Fluid Mech.*, **26** (1994), 191–210.
- [2] Ambrose, D. M., Well-posedness of vortex sheets with surface tension, preprint
- [3] Amick, C. J., Bounds for water waves. *Arch. Rat. Mech. Anal.*, **99** (1987), 99–114.
- [4] Amick, C. J. and K. Kirchgässner, A theory of solitary water-waves in the presence of surface tension. *Arch. Rat. Mech. Anal.*, **105** (1989), 1–49.
- [5] Amick, C. J. and J. F. Toland, On solitary water-waves of finite amplitude. *Arch. Rat. Mech. Appl.*, **76** (1981), 9–96.
- [6] Amick, C. J. and J. F. Toland, On periodic water-waves and their convergence to solitary waves in the long-wave limit. *Phil. Trans. Roy. Soc. London*, **A303** (1981), 633–673.
- [7] Amick, C. J. and J. F. Toland, A differential equation in the theory of resonant oscillations of water waves. *Proc. Roy. Soc. Edinburgh*, **114A**(1990), 15–26.
- [8] Andreev, V. K., Vortical perturbation of unsteady motion of fluid with free surface. *Zh. Prikl. Mekh. Tekh. Fiz.*, **5** (1975), 58–68. (Russian)
- [9] Andreev, V. K., Stability of the vortical perturbation of unsteady motion of planar layered ideal fluid with free surface. *Izv. Akad. Nauk SSSR Ser. Mekh. Zhid. Gaza*, **2** (1986), 15–21. (Russian)

- [10] Beale, J. T., The existence of solitary water waves. *Comm. Pure Appl. Math.*, **30** (1977), 373–389.
- [11] Beale, J. T. The existence of cnoidal water waves with surface tension, *J. Differential Equations*, **31** (1979), 230–263.
- [12] Beale, J. T., Water waves generated by a pressure disturbance on a steady stream. *Duke Math. J.*, **47** (1980), 297–323.
- [13] Beale, J. T., Exact solitary water waves with capillary ripples at infinity. *Comm. Pure Appl. Math.*, **44** (1991), 211–257.
- [14] Beale, J. T., A convergent boundary integral method for three-dimensional water waves. *Math. Comp.*, **70** (2001), 977–1029.
- [15] Beale, J. T., T. Y. Hou, and J. S. Lowengrub, Growth rates for the linearized motion of fluid interfaces away from equilibrium. *Comm. Pure Appl. Math.*, **46** (1993), 1269–1301.
- [16] Beale, J. T., T. Y. Hou, and J. S. Lowengrub, Convergence of a boundary integral method for water waves. *SIAM J. Numer. Anal.*, **33** (1996), 1797–1843.
- [17] Bimenov, M. A., Well-posedness of a linear problem on small perturbations of the motion of an ideal fluid with a free boundary in the class of smooth functions. (Russian) *Dinamika Sploshn. Sredy*, **102**(1991), 18–30.
- [18] Bimenov, M. A., The three-dimensional Cauchy-Poisson problem in classes of functions of finite smoothness. (Russian) *Dinamika Sploshn. Sredy*, **109**(1994), 65–78.
- [19] Christodoulou, D. and H. Lindblad, On the motion of the free surface of a liquid. *Comm. Pure Appl. Math.*, **53** (2000), 1536–1602.
- [20] Craig, W., An existence theory for water waves and the Boussinesq and Korteweg-de Vries scaling limits. *Commun. Partial Differ. Equations*, **10** (1985), 787–1003.
- [21] Craig, W. and P. Sternberg, Symmetry of free-surface flows. *Arch. Rat. Mech. Anal.*, **118** (1992), 1–36.
- [22] Crapper, G. D., An exact solution for progressive capillary waves of arbitrary amplitude. *J. Fluid Mech.*, **2** (1957), 532–540.
- [23] Dgaygui, K. and P. Joly, Absorbing boundary conditions for linear gravity waves. *SIAM J. Appl. Math.*, **54**(1994), 93–131.
- [24] Ebin, D. G., The equations of motion of a perfect fluid with free boundary are not well posed. *Commun. Partial Differ. Equations*, **12**, (1987), 1175–1201.

- [25] Ebin, D. G., Ill-posedness of the Rayleigh-Taylor and Helmholtz problems for incompressible fluids. *Comm. Partial Differential Equations*, **13** (1988), 1265–1295.
- [26] Epenberg, V. B., Planar stationary problem with free boundary and corner angle for Euler equation. (Russian) *Dinamika Sploshn. Sredy*, **14** (1973), 131–141.
- [27] Friedrichs K. O. and D. J. Hyers, The existence of solitary waves. *Comm. Pure Appl. Math.*, **7** (1954), 517–550.
- [28] Garipov, R. M., On the linear theory of gravity waves: the theory of existence and uniqueness. *Arch. Rat. Mech. Anal.*, **24** (1967), 352–362.
- [29] Gerber, R., Sur les solutions exactes des équations du mouvement avec surface libre d'un liquide pesant. *J. Math. Pures Appl.*, **34** (1955), 185–299.
- [30] Hărăguș, M., Model equations for water waves in the presence of surface tension. *Eur. J. Mech., B/Fluids*, **15** (1996), 471–492.
- [31] Hou, T. Y. and Pingwen Zhang, Growth rates for the linearized motion of 3-D fluid interfaces with surface tension far from equilibrium. *Asian J. Math.*, **2** (1998), 263–288.
- [32] Hou, T. Y., Zhen-huan, Teng and Pingwen Zhang, Well-posedness of linearized motion for 3-D water waves far from equilibrium. *Comm. Partial Differential Equations*, **21** (1996), 1551–1585.
- [33] Iguchi, T., Two-phase problem for two-dimensional water-waves of finite depth. *Math. Models Meth. Appl. Sci.*, **7** (1997), 791–821.
- [34] Iguchi, T., On the irrotational flow of incompressible ideal fluid in a circular domain with free surface. *Publ. RIMS, Kyoto Univ.*, **34** (1998), 525–565.
- [35] Iguchi, T., Well-posedness of the initial value problem for capillary-gravity waves. *Funccial. Ekvac.*, **44** (2001), 219–241.
- [36] Iguchi, T., On steady surface waves over a periodic bottom: relation between the pattern of imperfect bifurcation and the shape of the bottom. *Wave Motion*, **37** (2003), 219–239.
- [37] Iguchi, T., N. Tanaka and A. Tani, On the two-phase free boundary problem for two-dimensional water waves. *Math. Ann.*, **309** (1997), 199–223.
- [38] Iguchi, T., N. Tanaka and A. Tani, On a free boundary problem for an incompressible ideal fluid in two space dimensions. *Adv. Math. Sci. Appl.*, **9** (1999), 415–472.

- [39] Iooss, G. and K. Kirchgässner, Water waves for small surface tension: an approach via normal form. *Proc. Roy. Soc. Edinburgh*, **122A** (1992), 267–299.
- [40] Kano, T., Une théorie trois-dimensionnelle des ondes de surface de l'eau et le développement de Friedrichs. *J. Math. Kyoto Univ.*, **26** (1986), 101–155 and 157–175.
- [41] Kano, T. and T. Nishida, Sur les ondes de surface de l'eau avec une justification mathématique des équations des ondes en eau peu profonde. *J. Math. Kyoto Univ.*, **19** (1979), 335–370.
- [42] Kano, T. and T. Nishida, A mathematical justification for Korteweg-de Vries equation and Boussinesq equation of water surface waves. *Osaka J. Math.*, **23** (1986), 389–413.
- [43] Krasovskii, Ju. P., On the theory of steady-state waves of finite amplitude. *U.S.S.R. Comput. Math. and Math. Phys.*, **1** (1961), 996–1018.
- [44] Lamb, H., *Hydrodynamics*. 6th edition, Cambridge University Press, 1932.
- [45] Lavrentiev, M. A., On the theory of long waves. *Acad. Nauk Ukrain. R. S. R., Zbornik Prac. Inst. Mat.* V. 1946, No. 8 (1947), 13–69. [Ukrainian]
- [46] Lavrentiev, M. A., On some problems of motion of fluid with free surfaces. *Prikl. Mat. Mekh.*, **30** (1966), 177–182. [Russian]
- [47] Lavrentiev, M. A., To the theory of long waves. *Zhur. Prikl. Mekh. Tekh. Fiz.*, **5** (1975), 3–46. [Russian]
- [48] Levi-Civita, T., Détermination rigoureuse des ondes permanentes d'ampleur finie, *Math. Ann.*, **93** (1925), 264–314.
- [49] Lindblad, H., The motion of the free surface of a liquid. *Seminaire: Equations aux Derivées Partielles*, 2000–2001, Exp. No. VI, 10 pp., *Semin. Equ. Deriv. Partielles*, Ecole Polytech., Palaiseau, 2001.
- [50] Lindblad, H., Well-posedness for the linearized motion of an incompressible liquid with free surface boundary. *Comm. Pure Appl. Math.*, **56** (2003), 153–197.
- [51] Nalimov, V. I., A priori estimates of solutions of elliptic equations in the class of analytic functions and their applications to the Cauchy-Poisson problem. *Soviet Math. Dokl.*, **10** (1969), 1350–1354.
- [52] Nalimov, V. I., New model of the Cauchy-Poisson problem. *Dinamika Splošn. Sredy*, **12** (1972), 86–123. [Russian]

- [53] Nalimov, V. I., The Cauchy-Poisson problem. *Dinamika Splošn. Sredy*, **18** (1974), 104–210. [Russian]
- [54] Nalimov, V. I., Stationary surface waves over an uneven bottom. *Dinamika Splošn. Sredy*, **58** (1982), 108–155. [Russian]
- [55] Nalimov, V. I., Two problems in plane stationary surface wave theory. *Free boundary problems in fluid flow with applications (Montreal, PQ, 1990)*, 24–27, Pitman Res. Notes Math. Ser., **282**, Longman Sci. Tech., Harlow, 1993.
- [56] Nalimov, V. I., A model problem on vortex surface waves. *Sibirisk Mat. Zh.*, **35**(1994), 1119–1124; English Translation *Siberian Math. J.*, **35**(1996), 997–1001.
- [57] Nalimov, V. I., Unsteady vortex surface waves. *Sibirisk Mat. Zh.*, **37**(1996), 1356–1366; English Translation *Siberian Math. J.*, **37**(1996), 1189–1198.
- [58] Nalimov, V. I. and V. V. Pukhnachov, *non-stationary motion of ideal fluids with free boundary*. Novosibirsk State University, Novosibirsk, 1975. [Russian]
- [59] Nekrasov, A. I., On steady waves. *Izv. Ivanovo-Vosnosensk. Politehn. Inst.*, **3** (1921), 52–65. [Russian]
- [60] Nekrasov, A. I., *On steady waves, The exact theory of steady waves on the surface of a heavy fluid*. Izdat. Akad. Nauk SSSR, Moscow, 1951. [Russian]
- [61] Nishida, T., Analysis of fluid equations. Free surface problems. *Sûgaku*, **37** (1985), 289–304. [Japanese]
- [62] Nishida, T., Equations of fluid dynamics—free surface problems. Frontiers of the mathematical sciences: 1985. *Comm. Pure Appl. Math.*, **39** (1986), S221–S238.
- [63] Ogawa, M., Free surface motion of an incompressible ideal fluid. preprint.
- [64] Ogawa, M. and A. Tani, Free boundary problem for an incompressible ideal fluid with surface tension. *Math. Models Methods Appl. Sci.*, **12** (2002), 1725–1740.
- [65] Ogawa, M. and A. Tani, Incompressible perfect fluid motion with free boundary of finite depth. To appear in *Adv. Math. Sci. Appl.*
- [66] Okamoto, H., Stationary free boundary problems for circular flows with or without surface tension. *Lecture Notes in Num. Appl. Anal.*, **5** (1982), 233–251.
- [67] Okamoto, H., Nonstationary or stationary free boundary problems for perfect fluid with surface tension. *Lecture Notes in Num. Appl. Anal.*, **6** (1983), 143–154.

- [68] Okamoto, H., Bifurcation phenomena in a free boundary problem for a circulating flow with surface tension. *Math. Meth. Appl. Sci.*, **6** (1984), 215–233.
- [69] Okamoto, H., Nonstationary free boundary problem for perfect fluid with surface tension. *J. Math. Soc. Japan*, **38** (1986), 381–401.
- [70] Okamoto, H., Interfacial progressive water waves – a singularity–theoretic view. *Tôhoku Math. J.*, **49** (1997), 33–57.
- [71] Ovsjannikov, L. V., A singular operator in a scale of Banach spaces. *Soviet Math. Dokl.*, **6** (1965), 1025–1028.
- [72] Ovsjannikov, L. V., A nonlinear Cauchy problem in a scale of Banach spaces. *Soviet Math. Dokl.*, **12** (1971), 1497–1502.
- [73] Ovsjannikov, L. V., To the shallow water theory foundation. *Arch. Mech.*, **26** (1974), 407–422.
- [74] Ovsjannikov, L. V., *Group analysis of differential equations*. Nauka, 1978. [Russian]
- [75] Ovsjannikov, L. V., N. I. Makarenko, V. I. Nalimov, V. Yu. Lyapidenskii, P. I. Plotnikov, I. V. Sturova, V. I. Bukreev and V. A. Vladimirov, *Nonlinear problems in the theory of surface and internal waves*. Nauka, Sibirisk. Otdel. Novosibirsk, 1985. [Russian]
- [76] Plotnikov, P. I. and J. F. Toland, Nash-Moser theory for standing water waves. *Arch. Rat. Mech. Anal.*, **159** (2001), 1–83.
- [77] Reeder, J. and M. Shinbrot, The initial value problem for surface waves under gravity. II. The simplest 3-dimensional case. *Indiana Univ. Math. J.*, **25** (1976), 1049–1071.
- [78] Schneider, G. and C. E. Wayne, The long-wave limit for the water wave problem I. The case of zero surface tension. *Comm. Pure Appl. Math.*, **53** (2000), 1475–1535.
- [79] Schneider, G. and C. E. Wayne, The rigorous approximation of long-wavelength capillary-gravity waves. *Arch. Ration. Mech. Anal.*, **162** (2002), 247–285.
- [80] Schneider, G. and C. E. Wayne, Estimates for the three wave interaction of surface water waves. preprint.
- [81] Serrin, J., Mathematical principles of classical fluid mechanics. *Handbuph der Physik* **8/1** (1959), 125–263. Springer-Verlag
- [82] Shercliff, J. A., Simple rotational flows. *J. Fluid Mech.*, **82** (1977), 687–703.

- [83] Shinbrot, M., The initial value problem for surface waves under gravity. I. The simplest case. *Indiana Univ. Math. J.*, **25** (1976), 281–300.
- [84] Stoker, J. J., *Water waves*. Interscience Publ., New York, 1957.
- [85] Struik, D. J., Détermination rigoureuse des ondes irrotationnelles périodiques dans un canal à profondeur finie. *Math. Ann.*, **95** (1926), 595–634.
- [86] Sulem, C., P. L. Sulem, C. Bardos and U. Frisch, Finite time analyticity for two and three dimensional Kelvin-Helmholtz instability. *Commun. Math. Phys.*, **80** (1981), 485–516.
- [87] Sulem, C. and P. L. Sulem, Finite time analyticity for the two- and three-dimensional Rayleigh-Taylor instability. *Trans. Amer. Math. Soc.*, **287** (1985), 127–160.
- [88] Tanaka, M., The stability of steep gravity waves. *J. Phys. Soc. Japan*, **52** (1983), 3047–3055.
- [89] Tanaka, M., The stability of steep gravity waves. Part 2. *J. Fluid Mech.*, **156** (1985), 281–289.
- [90] Taylor, G. I., The instability of liquid surfaces when accelerated in a direction perpendicular to their planes I. *Proc. Roy. Soc. London, Ser. A.*, **201** (1950), 192–196.
- [91] Vanden-Broeck, J.-M., The influence of surface tension on cavitating flow past a curved obstacle. *J. Fluid Mech.*, **133** (1983), 255–264.
- [92] Ursell, F. The long-wave paradox in the theory of gravity waves. *Proc. Cambridge Philos. Soc.*, **49** (1953), 685–694.
- [93] Wehausen, J. W. and E. V. Laitone, Surface waves. *Handbuch der Physik* (Ed. S. Flügge), **9**, Springer-Verlag, Berlin, 1960.
- [94] Wu, S., Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.*, **130** (1997), 39–72.
- [95] Wu, S., Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Amer. Math. Soc.*, **12** (1999), 445–495.
- [96] Wu, S., Recent progress in mathematical analysis of vortex sheets. Proceedings of the International Congress of Mathematicians, Vol. III (Beijing, 2002), 233–242, Higher Ed. Press, Beijing, 2002.

- [97] Yosihara, H., Gravity waves on the free surface of an incompressible perfect fluid of finite depth. *Publ. RIMS Kyoto Univ.*, **18** (1982), 49–96.
- [98] Yosihara, H., Capillary-gravity waves for an incompressible ideal fluid. *J. Math. Kyoto Univ.*, **23** (1983), 649–694.
- [99] Zufiria, J. A., Symmetry breaking in periodic and solitary gravity–capillary waves on water of finite depth. *J. Fluid Mech.*, **184** (1987), 183–206.